

Self-avoiding Gonihedric String and Spin Systems

G.K. Savvidy

Physics Department, University of Crete
71409 Iraklion, Crete, Greece
and

Institut für Theoretische Physik
D-6000 Frankfurt am Main 11, Germany

K.G. Savvidy

Moscow Institute of Physics and Technology
141700 Moscow, Russia

Abstract

We classify different theories of self-intersecting random surfaces assigning special weights to intersections. When self-intersection coupling constant κ tends to zero, then the surface can freely intersect and it is completely self-avoiding when κ tends to infinity. Equivalent spin systems for this general case were constructed. In two-dimension the system with $\kappa = 0$ is in complete disorder as it is in the case of 2D gauge Ising system.

1.

In the recent article [1] the authors formulated gonihedric string, which was introduced in [2, 3, 4], on a euclidian lattice and have found such spin systems with local interaction which reproduce the same surface dynamics.

In the case of soft-self-avoiding surfaces they found, that the equivalent spin system coincides with usual or gauge Ising ferromagnet with an additional diagonal antiferromagnetic interaction. The coupling constants of this spin systems are specially adjusted. In the present article we shall consider the more general case of self-avoiding and non-self-avoiding surfaces.

In a continuum euclidean space R^d a random surface is associated with a connected polyhedral surface M with vertex coordinates X_i , where $i = 1, \dots, |M|$ and $|M|$ is the number of the vertices. The energy functional for the polyhedral surface is defined as [2]

$$A(M) = \sum_{\langle i,j \rangle} |X_i - X_j| \cdot \Theta(\alpha_{i,j}), \quad (1)$$

where

$$\begin{aligned} \Theta(\pi) &= 0, \\ \Theta(2\pi - \alpha) &= \Theta(\alpha), \\ \Theta(\alpha) &\geq 0, \end{aligned} \quad (2)$$

the summation is over all edges $\langle i, j \rangle$ of M and $\alpha_{i,j}$ is the angle between two neighbor faces of M in R^d having a common edge $\langle i, j \rangle$.

First we will classify different theories of self-intersecting surfaces assigning special weights to intersections [4]. For non-self-avoiding surfaces it is assumed that self-intersections do not produce any additional energy and thus the surface can freely intersect. In other cases self-intersections of a surface produce an additional energy and one should define nontrivial weights associated with these self-intersections. It was suggested in [4] to assign the following weights to intersection

$$\kappa \sum_{\text{common edges}} |X_i - X_j| \cdot (\Theta(\alpha_{i,j}^{(1)}) + \dots + \Theta(\alpha_{i,j}^{(r(2r-1))})), \quad (3)$$

where κ is self-intersection coupling constant, $\alpha_{i,j}^{(1)}, \dots, \alpha_{i,j}^{(r(2r-1))}$ are the angles between the pairs of faces of M at the common edge $\langle i, j \rangle$ in R^d in which $2r$ faces intersect $r = 0, 1, \dots, d-1$. It is natural to use κ for the definition of the different surface systems.

In euclidean space R^d the coupling constant κ can be fixed from physical requirement. Indeed from continuity principle [4] it follows that self-intersection coupling constant κ should be

$$\kappa = \frac{1}{2} \quad (4)$$

Physically this means that crumpled surface with folds can easily tear into peaces in the places where the surface has acute angles. Continuity principle guarantees that these transitions have non-zero amplitude even at the *classical level*.

On the lattice in Z^d the surface is associated with the collection of plaquettes, whose edgies are glued together pairwise and the surface is closed if each edge of the lattice belongs to an even number of plaquettes. The angles between plaquettes are either $0, \pi/2$ or π . There are no "continuous deformations" of the surface on the lattice with respect to angles. Therefore continuity principle does not tell us too much about the way how to choose κ on the lattice and we should keep κ as a free parameter of the theory.

Thus one can distinguish different cases through the value of κ . We will define three regions:

- i) $\kappa = 0$ - "bosonic" case
- ii) $0 < \kappa < \infty$ - "soft-fermionic" case
- iii) $\kappa = \infty$ - "fermionic" case.

In the last case the surface is completely self-avoiding. In [1] the authors considered the soft-fermionic case $\kappa = 1$.

Now we have a new set of rules (1) together with (3) for the energy functional and therefore can define the surface dynamics on the lattice with different self-intersection coupling constant κ . The energy of the lattice surface can be reexpressed through the plaquette variables U_P ($U_P = -1$ if $P \in M$ and $U_P = +1$ if $P \notin M$) [1]. The energies of all possible eight configurations at a given lattice edge in Z^3 can be expressed in the form

$$H_{edge}^\kappa = \frac{\kappa}{4}(2 - U_1 - U_{-1})(2 - U_2 - U_{-2}) + \frac{1 - \kappa}{4}(2 - U_1 U_{-1} - U_2 U_{-2}), \quad (5)$$

where the given edge is in the 3rd direction and since only an even number of plaquettes can meet at a given lattice edge, the total product of corresponding plaquette variables U_P surrounding the edge is always equal to one $U_1 U_{-1} U_2 U_{-2} = 1$. In (5) we have an extra term which is zero in the case $\kappa = 1$ considered in [1].

In four-dimension, Z^4 , again we have more complicated structure, than in the case $\kappa = 1$. We find that

$$\begin{aligned} H_{edge}^\kappa &= \frac{\kappa}{4}(2 - U_1 - U_{-1})(2 - U_2 - U_{-2}) \\ &\quad + \frac{\kappa}{4}(2 - U_2 - U_{-2})(2 - U_3 - U_{-3}) \\ &\quad + \frac{\kappa}{4}(2 - U_3 - U_{-3})(2 - U_1 - U_{-1}) \\ &\quad + \frac{1 - \kappa}{4} [(U_1 + U_{-1})(1 - U_2 U_{-2}) + (U_2 + U_{-2})(1 - U_3 U_{-3}) + (U_3 + U_{-3})(1 - U_1 U_{-1})]. \end{aligned} \quad (6)$$

The full energy associated with the lattice surface can always be written as

$$H^\kappa = \sum_{\text{over all edges}} H_{edge}^\kappa \quad (7)$$

Attaching spin variables to vertices of the dual lattice in Z^3 and to links in Z^4 one can get the Hamiltonian in terms of the independent spin variables $\sigma = \pm 1$ [5]. In Z^3 from (5) we get

$$\begin{aligned}
H^\kappa = & -2\kappa \sum_{\vec{r}, \vec{\alpha}} \sigma_{\vec{r}} \sigma_{\vec{r}+\vec{\alpha}} + \frac{1}{2}\kappa \sum_{\vec{r}, \vec{\alpha}, \vec{\beta}} (\sigma_{\vec{r}} \sigma_{\vec{r}+\vec{\alpha}+\vec{\beta}} + \sigma_{\vec{r}+\vec{\alpha}} \sigma_{\vec{r}+\vec{\beta}}) \\
& - \frac{1-\kappa}{2} \sum_{\vec{r}, \vec{\alpha}, \vec{\beta}} \sigma_{\vec{r}} \sigma_{\vec{r}+\vec{\alpha}} \sigma_{\vec{r}+\vec{\alpha}+\vec{\beta}} \sigma_{\vec{r}+\vec{\beta}}, \tag{8}
\end{aligned}$$

where $\vec{r}$ is a three-dimensional vector whose components are integer and $\vec{\alpha}, \vec{\beta}$ are unit vectors parallel to axis. The Hamiltonian (8) is invariant under global Z_2 symmetry. But as it is easy to see from (8) the ground state of the system has larger degeneracy than the ground state of the ferromagnet, parallel layers of differently oriented spins have the same energy. This has a strong influence on the phase structure of the system. High degeneracy of the ground state resembles gauge spin systems with local Z_2 symmetry [5].

In Z^4 from (6) we get

$$\begin{aligned}
H^\kappa = & -\frac{5\kappa-1}{g^2} \sum_{\text{plaquettes}} (\sigma\sigma\sigma\sigma) + \frac{\kappa}{4g^2} \cdot \sum_{\text{right angle plaquettes}} (\sigma\sigma\sigma\sigma_\alpha)^{rt} (\sigma_\alpha\sigma\sigma\sigma) \\
& - \frac{1-\kappa}{4g^2} \cdot \sum_{\text{triples of right angle plaquettes}} (\sigma\sigma\sigma\sigma_\alpha)^{rt} (\sigma_\alpha\sigma\sigma\sigma_\beta)^{rt} (\sigma_\beta\sigma\sigma\sigma), \tag{9}
\end{aligned}$$

As it is seen from (8) and (9) the difference with the case $\kappa = 1$ [1] consists in the fact that in (8) appears a new four-spin interaction term and a three-plaquette interaction term in (9). The partition function is defined as

$$Z(\beta) = \sum_{\{\sigma_{\vec{r}}\}} e^{-\beta H^\kappa}. \tag{10}$$

Expressions (5),(6) and (8),(9) together with (7) and (10) completely define the desired system of random surfaces on the lattice. The advantage of the given above formulation of the initial system of the random surfaces as a spin system with local interaction consists in the fact that now one can study more familiar spin systems and their phase structure.

2.

The systems of the same spirit can be formulated in any dimension [1]. Below we will concentrate on the two-dimensional case where they present a model of random walks [1].

The amplitudes of the elementary steps on the two-dimensional square lattice can be represented in terms of the eight-vertex model [6, 7, 8, 9] with the weights (see (5) with the correspondence: $3D$ edge $\longleftrightarrow$ $2D$ vertex, $3D$ plaquettes $\longleftrightarrow$ $2D$ bonds)

$$\omega_1 = 1, \quad \omega_2 = \exp(-4\beta\kappa), \quad \omega_3 = \omega_4 = 1, \quad \omega_5 = \omega_6 = \omega_7 = \omega_8 = \exp(-\beta), \tag{11}$$

where $\omega_\xi = \exp(-\beta\epsilon_\xi)$ and ϵ_ξ is the energy assigned to the ξ th type of vertex configuration ($\xi_i = 1, \dots, 8$)

$$\epsilon_1 = 0, \quad \epsilon_2 = 4\kappa, \quad \epsilon_3 = \epsilon_4 = 0, \quad \epsilon_5 = \epsilon_6 = \epsilon_7 = \epsilon_8 = 1. \quad (12)$$

The partition function (10) therefore becomes equal to (see (5) and (7))

$$Z(\omega_1, \dots, \omega_8) = \sum_{\text{over all closed paths}} \prod_{i=1}^{N^2} \omega_{\xi_i} \quad (13)$$

where the summation is extended over all closed paths consisting of vertices of even degrees (0, 2 or 4) on lattice with $N \times N$ sites. The evolution of the partition function (13) is a special case of more general problem involving the counting of weighted graphs (paths) with 16 kinds of vertex configurations with arbitrary weights [8]. In our case (11) $\omega_9 = \dots = \omega_{16} = 0$.

The partition function (13) possesses a number of evident symmetries [7]

$$\begin{aligned} Z(\omega_1, \omega_2, \omega_3, \omega_4; \omega_5, \omega_6, \omega_7, \omega_8) &= Z(\omega_2, \omega_1, \omega_4, \omega_3; \omega_6, \omega_5, \omega_8, \omega_7) \\ &= Z(\omega_4, \omega_3, \omega_2, \omega_1; \omega_7, \omega_8, \omega_5, \omega_6) = Z(\omega_3, \omega_4, \omega_1, \omega_2; \omega_8, \omega_7, \omega_6, \omega_5). \end{aligned} \quad (14)$$

Less evident is the duality transformation of Nagle [10, 8, 11]. We shall write here only the resulting linear transformation of the eight weights (11) into 16 kinds of vertex configurations appearing in the dual representation [8]

$$\omega_{\xi_i}^* = \sum_{\xi_i=1}^8 \omega_{\xi_i} \prod_k c_{i,k}(\xi_i), \quad (15)$$

where one should draw a bond between vertices i and k for each $c_{i,k}(\xi_i)$

$$c_{i,k}(\xi_i) = c_{k,i}(\xi_i) = \begin{cases} 1 & \text{if } \xi_i \text{ has bond on the edge } \langle i, k \rangle \\ -1 & \text{if } \xi_i \text{ does not have bond on the edge } \langle i, k \rangle \end{cases} \quad (16)$$

The summation in (13) should be extended now to all graphs which include all 16 kinds of vertices of even and odd degrees. It is easy to compute these dual weights from (15) and (11)

$$\begin{aligned} a^* &= \omega_1^* = \omega_2^* = 4(1 + \omega) - (1 - \omega^{4\kappa}); & b^* &= \omega_3^* = \omega_4^* = 4(1 - \omega) - (1 - \omega^{4\kappa}); \\ c^* &= \omega_5^* = \omega_6^* = 1 - \omega^{4\kappa}; & d^* &= \omega_7^* = \omega_8^* = 1 - \omega^{4\kappa}; & \omega_9^* &= \dots = \omega_{16}^* = 1 - \omega^{4\kappa}, \end{aligned} \quad (17)$$

where $\omega = \exp(-\beta)$. In the bosonic case i) $\kappa = 0$ the dual system (17) reduces to the simple eight-vertex model

$$a^* = 1 + \omega, \quad b^* = 1 - \omega, \quad c^* = d^* = 0 \quad (18)$$

which can be easily solved [9]. The free energy f per vertex is

$$-\beta f = \ln a^* = \ln \frac{a + b + c + d}{2} = \ln(1 + \omega), \quad (19)$$

where $a = \omega_1 = \omega_2 = 1$, $b = \omega_3 = \omega_4 = 1$, $c = \omega_5 = \omega_6 = \omega$, $d = \omega_7 = \omega_8 = \omega$. The free energy (19) is an analytic function of β and the system is completely disordered (dual model (18) is in ferroelectric order $a^* > b^* + c^* + d^*$). This is a result of the high degeneracy of the ground state: the vertices ω_1 , ω_2 , ω_3 and ω_4 have the same energy and the ground state is no longer unique and has high degeneracy. The transition temperature vanishes as in the case of the gauge Ising model in two-dimension [5] or Ising model in one-dimension.

There are two large classes of the general eight-vertex model which can be solved analitically: α) free-fermion model [12, 13, 14, 7, 9], when

$$\omega_1\omega_2 + \omega_3\omega_4 = \omega_5\omega_6 + \omega_7\omega_8 \quad (20)$$

and β) zero-field eight-vertex Baxter model [6], when $a = \omega_1 = \omega_2$, $b = \omega_3 = \omega_4$, $c = \omega_5 = \omega_6$, $d = \omega_7 = \omega_8$. Nor of these conditions is fulfilled by the weights (11), except of the case i) $k = 0$, which was already considered (18), (19).

To get information about the behavior of the general system (11), when $k \neq 0$, one can use fermionic representation of the partition function (13). Following [12, 13, 14, 7, 9] one can rewrite the partition function (13) as a product of the fermion operators

$$\begin{aligned} Z(\omega_1, \dots, \omega_8) = < 0 | \prod_{i=1}^{N^2} (\omega_1 + \omega_2 a_i^+ b_i^+ a_{i-N} b_{i-1} + \omega_3 a_i^+ a_{i-N} + \omega_4 b_i^+ b_{i-1} + \\ + \omega_5 a_i^+ b_{i-1} + \omega_6 b_i^+ a_{i-N} + \omega_7 a_i^+ b_i^+ + \omega_8 a_{i-N} b_{i-1}) | 0 >, \end{aligned} \quad (21)$$

where a_i^+ , b_i^+ and a_i , b_i are bond creation and annihilation fermion operators at the i 'th vertex, a refers to bonds in the vertical direction, b to bonds in the horizontal direction and $|0 >$ is the vacuum state. Using anticommutation relations of the fermion operators one can get [12, 13, 14, 7, 9]

$$Z = \omega_1^{N^2} < 0 | T \exp \left\{ \sum_{i=1}^{N^2} H(i) + g \cdot H_{int}(i) \right\} | 0 >, \quad (22)$$

where $H(i)$ is quadratic in the field operators

$$\omega_1 H(i) = \omega_3 a_i^+ a_{i-N} + \omega_4 b_i^+ b_{i-1} + \omega_5 a_i^+ b_{i-1} + \omega_6 b_i^+ a_{i-N} + \omega_7 a_i^+ b_i^+ + \omega_8 a_{i-N} b_{i-1}, \quad (23a)$$

$H_{int}(i)$ is quartic

$$H_{int}(i) = a_i^+ b_i^+ a_{i-N} b_{i-1} \quad (23b)$$

and

$$g = \frac{\omega_1\omega_2 + \omega_3\omega_4 - \omega_5\omega_6 - \omega_7\omega_8}{\omega_1^2} \quad (23c)$$

and T orders the indices from $i = 1$ to $i = N^2$.

For the system (11) quadratic part is equal to

$$H(i) = a_i^+ a_{i-N} + b_i^+ b_{i-1} + \omega(a_i^+ b_{i-1} + b_i^+ a_{i-N} + a_i^+ b_i^+ + a_{i-N} b_{i-1}) \quad (24a)$$

and the coupling constant (23c) is

$$g = 1 - 2\omega^2 + \omega^{4\kappa} \quad (24c)$$

One can directly use field theoretical methods to evaluate the expression (22). In high temperature regim when $\beta \rightarrow 0$ the coupling constant (24c) is small $g = 4(1 - \kappa)\beta + 4(2\kappa^2 - 1)\beta^2$, and one can use perturbation expansion. In zero approximation we have

$$\begin{aligned} -\beta f_0 &= \frac{1}{N^2} \ln Z_0 = \frac{1}{N^2} \ln \langle 0 | \text{Temp} \{ \sum_{i=1}^{N^2} H(i) \} | 0 \rangle \\ &= \frac{1}{8\pi^2} \int_0^{2\pi} \ln[(1 - \omega^2 + \omega^4) + (1 - \omega^2)(\cos \theta + \cos \theta \cos \phi + \cos \phi)] d\theta d\phi \end{aligned} \quad (25)$$

The free energy is analitic for all temperatures β and system is always in disordered regime. The physical reason of this type of phase structure of the system lies in the fact, that in (11) we don't have dominant vertex, the nessesary requirment for the order at low temperatures. Of course this is a zero order result and as it is easy to see from (24a) and (21) the expression (25) is reliable only if $2\omega^2 - 1 > 0$, that is for temperatures T greater than $\frac{1}{\ln \sqrt{2}}$. It is also true that at low tempetatures the coupling constant (24c) grows and is of order one. Therefore more rigor analysis should be done in the low temperature region to confirm zero order result [15, 16].

We expect more complicated phase structure of the system in high dimensions where the energy functional has the meaning of the size of the system and where it is not conformal invariant, as it is in the two-dimension case (11). Some exact results concerning three-dimensional edge models have been obtained in [17, 18, 19, 20]

In conclusion we would like to stress that (9) can be considered as an extention of the Wegner-Wilson action if one exchange spin variables σ by the elements $U = \exp(igA)$ of $SU(N)$. The new action will contain high derivative terms with specially adjusted coupling constants. This type of extension has been considered before in [21] with the aim to improve continuum limit of the lattice gauge theory.

Acknowledgements

One of the authors (G.K.S.) is grateful to Alexander Belavin for the discussion of the Wegner spin systems which took place in Yerevan in 1978 and to Franz Wegner for numerous stimulating conversations and kind hospitality in Heidelberg. We thank G.Athanasiu, E. Floratos, R. Flume, N. Papanicolaou, E. Paschos for interesting discussions and W. Greiner for useful comments and hospitality in Frankfurt University. One of the authors (K.G.S.) is thankful to S.Pnevmatikos for kind invitation into Summer School of Heraklion.

This work was supported in part by the Alexander von Humboldt Foundation.

References

- [1] G.K.Savvidy and F.J.Wegner. Geometrical string and spin systems. Preprint Crete-TH-5, July 1993; hep-th 9308094; Nucl.Phys.B(1993).
- [2] R.V. Ambartzumian, G.K. Savvidy, K.G. Savvidy and G.S. Sukiasian. Phys. Lett. B275 (1992) 99
- [3] G.K. Savvidy and K.G. Savvidy. String fine tuning. Preprint UFTP 302/1992; hep-th 9208041; Int. J. Mod. Phys. A (1993)
- [4] G.K. Savvidy, K.G. Savvidy. Gonihedric String and Asymptotic Freedom. Preprint UFTP 320/1992; hep-th 9301001; Mod.Phys.Lett.A (1993)
- [5] F.J. Wegner, J. Math. Phys.12 (1971) 2259
- [6] R.J.Baxter, Exactly solved models in statistical mechanics. Academic Press, London 1982
- [7] C.Fan and F.Y.Wu. Phys.Rev.179 (1969)560
- [8] F.Y.Wu. Phys.Rev.183 (1969) 604
- [9] C.Fan and F.Y.Wu. Phys.Rev.B2 (1970) 723
- [10] J.F.Nagle. J.Math.Phys.9 (1968) 1007
- [11] F.J. Wegner, Physica 68 (1973) 570
- [12] C.A.Hurst and H.S.Green. J.Chem.Phys. 33(1960) 1059
- [13] C.A.Hurst J.Math.Phys.7 (1966)305
- [14] R.W.Gibberd and C.A.Hurst J.Math.Phys. 8 (1967) 1427
- [15] R.E.Peierls. Proc. Cambridge Phil. Soc. 32 (1936) 477.
- [16] E.I. Dinaburg and Ya.G. Sinai, Comm.Math.Phys. 98 (1985) 119
- [17] J.F.Nagle, Comm.Math.Phys. 13 (1969) 62
- [18] C.Thibaudier and J.Villain, J.Phys.C: Solid State Phys. 5 (1972) 3429
- [19] P.Orland, Int. J. Mod.Phys. 5 (1991) 2401
- [20] N.Papanicolaou, Nucl.Phys. B305 (1988) 367
- [21] P.Weisz, Nucl.Phys. B212 (1983) 1